

REVIEW

Understanding of COVID-19 based on current evidence

Pengfei Sun¹ | Xiaosheng Lu² | Chao Xu¹ | Wenjuan Sun³ | Bo Pan⁴ 

¹Department of Plastic Surgery, Zibo Central Hospital, Zibo, China

²Department of Plastic Surgery, Affiliated Hospital of Weifang Medical University, Weifang, China

³Department of Medicine, People's Hospital of Boxing, Binzhou, China

⁴Department of Plastic Surgery, Plastic Surgery Hospital, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, China

Correspondence

Bo Pan, MD, Department of Plastic Surgery, Plastic Surgery Hospital, Peking Union Medical College, Chinese Academy of Medical Sciences, Bada Rd, Shijingshan, Beijing 100144, China.
Email: zbzbzhc@163.com

Abstract

Since December 2019, a series of unexplained pneumonia cases have been reported in Wuhan, China. On 12 January 2020, the World Health Organization (WHO) temporarily named this new virus as the 2019 novel coronavirus (2019-nCoV). On 11 February 2020, the WHO officially named the disease caused by the 2019-nCoV as coronavirus disease (COVID-19). The COVID-19 epidemic is spreading all over the world, especially in China. Based on the published evidence, we systematically discuss the characteristics of COVID-19 in the hope of providing a reference for future studies and help for the prevention and control of the COVID-19 epidemic.

KEYWORDS

2019-nCoV, coronavirus, COVID-19, epidemiology, pneumonia, SARS-CoV-2

1 | INTRODUCTION

Since December 2019, there has been a series of unexplained cases of pneumonia reported in Wuhan, China. The Chinese government and researchers took rapid measures to control the epidemic and carried out etiological research. On 12 January 2020, the World Health Organization (WHO) tentatively named this new virus as the 2019 novel coronavirus (2019-nCoV). On 30 January 2020, WHO announced the 2019-nCoV epidemic a public health emergency of international concern. On 11 February 2020, the WHO formally named the disease triggered by 2019-nCoV as coronavirus disease 2019 (COVID-19). On the same day, the coronavirus study group of the International Committee on Taxonomy of Viruses named 2019-nCoV as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). On 23 February 2020, there were 77 041 confirmed cases of SARS-CoV-2 infection in China. The number of infections has exceeded that of the SARS outbreak in China in 2002.^{1,2} To help healthcare workers around the world to better deal with the SARS-CoV-2, we review the relevant published papers on COVID-19 to provide a reference for future COVID-19 research.

1.1 | Sources of SARS-CoV-2

SARS-CoV-2 is a coronavirus and belongs to the β -coronavirus cluster. COVID-19 is the third known zoonotic coronavirus disease after SARS and the Middle East respiratory syndrome (MERS). SARS-CoV and MERS-CoV also belong to the β -coronavirus cluster.³ Zhu et al⁴ confirmed that SARS-CoV-2 was a new β -coronavirus belonging to the subgenus botulinum of Coronaviridae. According to the current data,⁵ the early COVID-19 cases were related to the Huanan seafood market, and the possibility of human-to-human transmissions could not be ruled out. The WHO report claimed that the SARS-CoV-2 could be detected in the environmental samples collected from the seafood market, but it has not yet been determined if a specific animal species carries the SARS-CoV-2. A study by Ji et al⁶ showed that the SARS-CoV-2 was a chimeric virus between a bat coronavirus and a coronavirus of unknown origin. By comparing with other animals, they found that snakes are the most likely wildlife repository for the SARS-CoV-2.⁶ The research by Benvenuto et al⁷ showed that the SARS-CoV-2 was only closely related to the coronavirus isolated from Chinese chrysanthemum-headed bats in 2015. Their research supported the theory that the transmission chain started from bats to humans.

Chan et al⁸ and Hui et al⁹ confirmed that SARS-CoV-2 was a new coronavirus closely related to the bat SARS-CoV. Recently, Zhou et al¹⁰ and Wu et al¹¹ found that the sequence homology between SARS-CoV-2 and SARS-CoV was 79.5%. They also found that the SARS-CoV-2 had high homology with bat coronaviruses. Therefore, the current evidence strongly supports that the SARS-CoV-2 was derived from bats, although the intermediate hosts of SARS-CoV-2 remain to be determined.

1.2 | Epidemiological characteristics of COVID-19

A study by Wang et al¹² showed that from January 10 to 24, 2020, the number of people diagnosed with the SARS-CoV-2 infection in China increased by 31.4 times. On 23 February 2020, the number of people diagnosed with COVID-19 in China was 1879 times of that on 10 January 2020. They estimated the case fatality rate of COVID-19 to be 2.84% based on their patient pool. The authors also found that the ratio of male to female deaths was 3.25:1, the median age of death was 75 years, the median time from the first symptom to death was 14 days, and the median time from early symptoms to death in people aged 70 or older (11.5 days) was shorter than that in people under 70 years old (20 days). These findings suggest the disease may progress faster in the elderly than in the young.

A study by Li et al⁵ showed the median age of 425 patients infected with SARS-CoV-2 was 59 years, of which 56% were males, the average incubation period was 5.2 days, and almost half of the adult patients were 60 years old or older. In the early stages, the number of infected patients doubled every 7.4 days. The transmission rate of individual infected patients was 2.2. Although 55% of the earliest SARS-CoV-2-infected patients related to the Huanan seafood market, the number of unrelated cases have increased exponentially since late December 2019.

Huang et al¹³ showed among the 41 patients with SARS-CoV-2 infections in their study, 73% of the patients were male, and 32% of the patients had underlying diseases, including diabetes (eight patients), hypertension (six patients), and cardiovascular disease (six patients). The median age was 49 years. Out of the 41 patients, 27 patients were associated with the Huanan seafood market. The case fatality rate among the SARS-CoV-2-infected patients in this study was 15%. A study by Wu et al¹⁴ estimated the transmission rate of individual infected patients to be 0.3. The case fatality rate among the SARS-CoV-2-infected patients in this study was 14%.

1.3 | Mechanism, symptoms, and diagnosis of COVID-19

Zhao et al¹⁵ found that angiotensin-converting enzyme 2 (ACE2) was the receptor for SARS-CoV-2. In the normal human lung, ACE2 is expressed on type I and II alveolar epithelial cells. Among them, 83% of the type II alveolar cells have ACE2 expression. Men had a higher ACE2 level in their alveolar cells than women. Asians have a higher level of ACE2 expression in their alveolar cells than the White and

African American populations. The binding of SARS-CoV-2 on ACE2 causes an elevated expression of ACE2, which can lead to damages on alveolar cells. Damages to alveolar cells can, in turn, trigger a series of systemic reactions and even death. They also confirmed that Asian males are more susceptible to SARS-CoV-2 infection. Wrapp et al¹⁶ found that the receptor-binding ability of SARS-CoV-2 is 10 to 20 times stronger than that of SARS-CoV.

Huang et al¹³ found that 98% of the patients in their study had fevers, of which 78% had a temperature higher than 38°C. They reported that 76% of the patients had coughs, 44% of patients experienced fatigue and muscle pain, and 55% of patients had dyspnea. A small number of patients also developed expectoration (28%), headaches (8%), hemoptysis (5%), and diarrhea (3%). Laboratory tests found that 25% of infected patients had leukopenia and 63% had lymphocytopenia. The level of aspartate aminotransferase was elevated in 37% of the patients. Myocarditis was diagnosed in 12% of the patients, and the level of hypersensitive troponin I was significantly increased in these patients. Abnormalities in chest computed tomography (CT) images were found in 100% of the patients. Grinding glass-like and consolidation areas were found in 98% of the infected patients' bilateral lungs.

Zhu et al⁴ reported three cases of COVID-19. Patient 1 was a 49-year-old female with a fever (body temperature 37 to 38°C) and had coughs accompanied by chest discomfort. Four days after the onset of the disease, her coughs and chest discomfort aggravated, but her fever subsided. Patient 2, a 61-year-old male, also developed fevers and coughs at the initial stage of the disease. Respiratory distress appeared 7 days after the onset of symptoms and worsened within the next 2 days. He was treated with mechanical ventilation. Patient 3 was a 32-year-old male whose symptoms were not described in the article. Patients 1 and 3 recovered and were discharged from hospital after treatment, but patient 2 died after 20 days of treatment.

Guan et al¹⁷ reported 1099 cases of 2019-nCoV infection. They found that fevers (87.9%) and coughs (67.7%) were the most common symptoms. Diarrhea (3.7%) and vomiting (5.0%) were rare. Abnormalities in chest CT images were found in 96% of the SARS-CoV-2-infected patients, and lymphopenia was observed in 82.1% of them.

1.4 | Prevention and treatment of COVID-19

As of 23 February 2020, no COVID-19 vaccine has been successfully developed. At present, the treatments of patients with SARS-CoV-2 infection are mainly symptomatic treatments. The study by Huang et al¹³ reported that the most common complications in patients with 2019-nCoV infection were acute respiratory distress syndrome, followed by anemia, acute heart injuries, and secondary infections. Therefore, empirical antibiotics, antiviral therapy (oseltamivir), and systemic corticosteroids were often used for treatments. Patients with intractable hypoxemia were given invasive mechanical ventilation. Holshue et al¹⁸ used remdesivir in the treatment of patients with SARS-CoV-2 infection and achieved good results. Lu¹⁹ postulated that, in addition to antiviral interferers and antibiotics, neuraminidase

inhibitors, RNA synthesis inhibitors, and Chinese traditional medicine could also be used in the treatment of COVID-19. Nevertheless, the efficacy of these drugs still needs to be verified by clinical trials.

In the absence of effective treatments, the best way to deal with the SARS-CoV-2 epidemic is to control the sources of infection. Strategies include early diagnoses, reporting, isolation, and supportive treatments; timely release of epidemic information; and maintenance of social orders. For individuals, protective measures, including improving personal hygiene, wearing medical masks, adequate rest, and keeping rooms well ventilated, can effectively prevent SARS-CoV-2 infection.¹⁷

2 | DISCUSSION

Regarding the case fatality and transmission rates among patients with SARS-CoV-2 infection, the findings from various studies were different. The studies of Wang et al¹² showed that the case fatality rate was 2.84%. According to Huang et al,¹³ the number was 15%. The study by Wu et al¹⁴ estimated that the case fatality rate was 14%. In the study by Zhu et al,⁴ the number was 33%. Guan et al¹⁷ reported that the case fatality rate was 1.36%. According to the official data released by China, the case fatality of COVID-19 patients was 3.17%.² In terms of the transmission rate, Li et al⁵ showed that the transmission rate was 2.2 per patient. Wu et al¹⁴ estimated that the transmission rate was 0.3 per patient. Yang et al²⁰ reported that the transmission rate was 3.77. We think that the transmission rate of 2.2 per patient seems to be more in line with the current situation.² Different sample sizes and possible viral variations may have contributed to the differences between the studies. Studies with a larger sample size may give a better estimation of the case fatality rate and transmission rate of the COVID-19. We know that the other two major zoonotic coronavirus diseases, SARS and MERS which caused widespread transmission have case fatality rates of 9.6% and 35%, respectively.⁹ According to the official Chinese data, the case fatality rate among the SARS-CoV-2-infected patients was much lower than that of SARS and MERS.²

In response to the COVID-19 epidemic, we believe that the focus of future studies will still be on the development of COVID-19 vaccines and effective drugs to treat COVID-19. These studies will help further reduce the case fatality and transmission rates among SARS-CoV-2-infected patients. Moreover, super-spreaders were reported during the SARS and MERS epidemics.²¹ Although the transmission rate for SARS-CoV-2 patients is about 2.2 at present, the number of cases inside and outside Wuhan is increasing rapidly. With the progress of diagnostic technology, potential super-spreaders may be discovered in the future. In the prevention of the spread of SARS-CoV-2, asymptomatic spreaders also need to be focused.²²

In addition, Sheng et al²³ found that viral infections can increase the risk of pulmonary fibrosis. Therefore, pulmonary fibrosis may be one of the severe complications after patients recover from 2019-nCoV infections. The prevention of pulmonary fibrosis in patients recovered from 2019-nCoV infections is an issue that urgently needs to be addressed.

ACKNOWLEDGMENT

The authors acknowledge TopEdit LLC for linguistic editing and proofreading during the preparation of this manuscript.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

AUTHOR CONTRIBUTIONS

BP and PS had the idea for and designed the study and take responsibility for the integrity of the data and the accuracy of the data analysis. PS and XL contributed to the writing of the report. BP contributed to the critical revision of the report. CX and WS contributed to the statistical analysis. All authors contributed to data acquisition, data analysis, or data interpretation, and reviewed and approved the final version.

ORCID

Bo Pan  <http://orcid.org/0000-0001-8900-8479>

REFERENCES

- Gorbalenya AE, Baker SC, Baric RS, et al. Severe acute respiratory syndrome-related coronavirus: The species and its viruses—a statement of the coronavirus study group. *bioRxiv*. 2020. <https://doi.org/10.1101/2020.02.07.937862>
- National Health Commission's briefing on the pneumonia epidemic situation (in Chinese). 2020. <http://www.nhc.gov.cn/yjb/s7860/202001/9614b05a8cac4ffabac10c4502fe517c.shtml>. Accessed 23 February 2020.
- Chen Y, Liu Q, Guo D. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J Med Virol*. 2020;92:418-423. <https://doi.org/10.1002/jmv.25681>
- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382:727-733. <https://doi.org/10.1056/NEJMoa2001017>
- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020; NEJMoa2001316. <https://doi.org/10.1056/NEJMoa2001316>
- Ji W, Wang W, Zhao X, Zai J, Li X. Homologous recombination within the spike glycoprotein of the newly identified coronavirus may boost cross-species transmission from snake to human. *J Med Virol*. 2020;92: 433-440. <https://doi.org/10.1002/jmv.25682>
- Benvenuto D, Giovannetti M, Ciccozzi A, Spoto S, Angeletti S, Ciccozzi M. The 2019-new coronavirus epidemic: evidence for virus evolution. *J Med Virol*. 2020. <https://doi.org/10.1002/jmv.25688>
- Chan JFW, Kok KH, Zhu Z, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect*. 2020;9(1):221-236. <https://doi.org/10.1080/22221751.2020.1719902>
- Hui DS, I Azhar E, Madani TA, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health—the latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis*. 2020;91:264-266.
- Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020. <https://doi.org/10.1038/s41586-020-2012-7>
- Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature*. 2020. <https://doi.org/10.1038/s41586-020-2008-3>
- Wang W, Tang J, Wei F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol*. 2020. <https://doi.org/10.1002/jmv.25689>

13. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
14. Wu P, Hao X, Lau EHY, et al. Real-time tentative assessment of the epidemiological characteristics of novel coronavirus infections in Wuhan, China, as at 22 January 2020. *Euro Surveill*. 2020;25(3).
15. Zhao Y, Zhao Z, Wang Y, et al. Single-cell RNA expression profiling of ACE2, the putative receptor of Wuhan 2019-nCoV. *bioRxiv*. 2020. <https://doi.org/10.1101/2020.01.26.919985>
16. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the SARS-CoV-2 spike in the prefusion conformation. *medRxiv*. 2020. <https://doi.org/10.1101/2020.02.11.944462>
17. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *medRxiv*. 2020. <https://doi.org/10.1101/2020.02.06.20020974>
18. Holshue ML, DeBolt C, Lindquist S, et al. First case of 2019 novel coronavirus in the United States. *N Engl J Med*. 2020:NEJMoa2001191. <https://doi.org/10.1056/NEJMoa2001191>
19. Lu H. Drug treatment options for the 2019-new coronavirus (2019-nCoV). *Biosci Trends*. 2020. <https://doi.org/10.5582/bst.2020.01020>
20. Yang Y, Lu Q-B, Liu M-J, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. *medRxiv*. 2020. <https://doi.org/10.1101/2020.02.10.20021675>
21. Wong G, Liu W, Liu Y, Zhou B, Bi Y, Gao GF. MERS, SARS, and Ebola: the role of super-spreaders in infectious disease. *Cell Host Microbe*. 2015;18(4):398-401.
22. Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med*. 2020:NEJMc2001468. <https://doi.org/10.1056/NEJMc2001468>
23. Sheng G, Chen P, Wei Y, et al. Viral infection increases the risk of idiopathic pulmonary fibrosis: a meta-analysis. *Chest*. 2019. <https://doi.org/10.1016/j.chest.2019.10.032>

How to cite this article: Sun P, Lu X, Xu C, Sun W, Pan B. Understanding of COVID-19 based on current evidence. *J Med Virol*. 2020;1-4. <https://doi.org/10.1002/jmv.25722>